arXiv:0802.3743v3 [nucl-th] 14 Dec 2008

Study of Chemical Equilibration of a Baryon Rich QGP

Abhijit Sen
Lecturer, Department of Physics, Suri Vidyasagar College, Suri-731101, INDIA

Keywords: Parton, equilibration, Baryon Number, Chemical Potential,flavour change

PACS No. 12.38.Mh(Quark Gluon Plasma), 21.65(Quark Matter),25.75(Relativistic heavy ion collisions)

December 14, 2008

Abstract

Parton equilibration studies for a thermally equilibrated but chemically non-equilibrated quark-gluon plasma (QGP) , likely to be formed at the relativistic colliders at BNL and CERN is presented. Parton equilibration is studied enforcing baryon number conservation. Process like quark - flavour interchanging is also taken into consideration. The degree of equilibration is studied comparatively for the various reactions / constraints that are being considered.

1 Introduction

The parton plasma likely to be created in high energy heavy ion collisions is sure to be in a state of both thermal and chemical non-equilibria.Once thermal equilibrium is attained, the subsequent evolution takes place according to the laws of hydrodynamics.

It is known that [1] the initial system produced at RHIC energies have a finite non- zero baryon density. The anti-particle to particle ratios at mid-rapidity for $\sqrt{s_{NN}}$=130 GeV Au+Au collisions at the STAR collaboration,BNL report that there is a noticiable excess of baryons as compared to anti-baryons as reflected by the yields of $\frac{\overline{p}}{p}$ (hovering between 0.6 and 0.8), $\frac{\overline{\Lambda}}{\Lambda}$ (hovering between 0.7 and 0.8) [2] for varying transverse momentum, centrality and rapidity. In a subsequent reporting of the BRAHMS collaboration of BNL[3] it is seen that during Pb+Pb collisions at $\sqrt{s_{NN}}$= 200 GeV the observed $\frac{\overline{p}}{p}$ ratio is a maximum of around 0.8 at zero rapidity and falls steadily for higher rapidity values.

This excess of baryons (and hence quarks) over anti-baryons (and hence anti-quarks) clearly necessiates the inclusion of a chemical potential into the theoretical framework . Moreover, strong interaction conserves baryon number, which also must be incorporated in our scheme. The present piece of work is aimed at attaining such an objective.

2 Brief Review of Previous Works

.

Inclusion of chemical potential into the framework of Parton equilibration studies has been rather recent. Although in 1986 Matsui et.al reported [8] on strangeness production rates at nonzero chemical potential, it was not until the very end of the past century that a group from BARC,Mumbai, India included a chemical potential into chemical equilibration studies [4,5]. They considered a chemical potential (for massless quarks) that equalled the system temperature at all temperatures which was not realistic as it did not consider baryon number conservation. Furthermore, they did not distinguish between quarks and anti-quarks. The same criticism regarding chemical potential and baryon number conservation also applies to an attempt [6] (incomplete) by the present author during the preliminary stages of the current work. However, explicit masses and distinction between quarks and anti-quarks had been incorporated in that conference report. In 2004, He et.al. [7] undertook a much more complete study including baryon number conservation. They used an expansion of the number densities over powers of chemical potential. We basically follow the same line with a few points in difference with the work by He et.al.The points of difference are:

i) We explicitly introduce the strange quark mass

ii) We treat quarks and antiquarks separately.It can be noted here that although He et. al. [7] started off with quark-antiquark distinction, subsequently that distinction was put away.

iii) We include results of full phase space calculations for both quark-antiquark pair production and quark flavour changing processes.

3 The Chemical Potential

At this point let us note the following points:

1. The initial strangeness content of the QGP fireball is zero and moreover strangeness is very nearly conserved in strong interaction which clearly indicates that the strange quark chemical potential must be zero.

2. If we consider a pair production process in QGP (production of non-strange quark- antiquark pair) it can be argued that the net chemical potential on both sides of the reaction must be the same. Since gluons have no chemical potential clearly this would imply

$$\mu_q = -\mu_{\overline{q}}$$

...(1)

3. We treat the up and down flavours in the same footing and treat them as massless.

4. For a given initial baryon number density and initial values of the light quark & antiquark non-equilibrium fugacities (which is defined as the ratio of the number density to the equilibrium number density) we can find the initial light quark chemical potential by method of iteration.

5. The rate of change of chemical potential with time is dictated by the baryon number conservation equation.

4 Thermodynamic & Statistical Mechanic Treatments

4.1 Distribution functions

For a thermally equilibrated but chemically non-equilibrated parton plasma, the distribution functions are given by the Juttner distribution functions [9]. In the present paper, initially we use the full distributions. In the latter stage we shall adopt certain approximations in order to achieve some mathematical simplicity as also to confront some mathematical unsolvability that we shall encounter. The full Juttner distribution functions are

$$\begin{aligned} f_g &= \frac{\lambda_g}{e^{\varepsilon/T} - \lambda_g} \\ f_{q(\overline{q})} &= \frac{\lambda_{q(\overline{q})}}{e^{(\varepsilon \mp \mu_q)/T} + \lambda_{q(\overline{q})}} \qquad ...(2) \\ f_s = f_{\overline{s}} &= \frac{\lambda_s}{e^{\varepsilon/T} + \lambda_s} \end{aligned}$$

4.2 Basic Quantities

Using standard techniques, we can obtain the various thermodynamic quantities that are required to specify the state of the system. We obtain:

$$Q = Q_g + \Sigma_{i=u,d,s}(Q_i + Q_{\overline{i}}) \ = Q_g + 2 * (Q_q + Q_{\overline{q}}) + 2Q_s \qquad ...(3)$$

Q being the number density, energy density or pressure in general and q denoting the massless quark flavours.

This gives

$$\begin{aligned} n = \{ &\frac{16T^3}{\pi^2}\Sigma_{k=1}^{\infty}\frac{\lambda_g^k}{k^3} + \frac{12T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} + \lambda_{\overline{q}}^k e^{-\frac{k\mu_q}{T}}}{k^3}\} + \\ &\frac{6m_{0s}^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\lambda_s^k\frac{K_2(kx_s)}{(kx_s)}\} \end{aligned}$$

$$\begin{aligned} \varepsilon = \{ &\frac{48T^4}{\pi^2}\Sigma_{k=1}^{\infty}\frac{\lambda_g^k}{k^4} + \frac{36T^4}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} + \lambda_{\overline{q}}^k e^{-\frac{k\mu_q}{T}}}{k^4}\} + \\ &\frac{6m_{0s}^4}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\lambda_s^k\{\frac{3K_2(kx_s)}{(kx_s)^2} + \frac{K_1(kx_s)}{(kx_s)}\}\} \end{aligned}$$

and

$$\begin{aligned} p = \{ &\frac{16T^4}{\pi^2}\Sigma_{k=1}^{\infty}\frac{\lambda_g^k}{k^4} + \frac{12T^4}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} + \lambda_{\overline{q}}^k e^{-\frac{k\mu_q}{T}}}{k^4}\} + \\ &\frac{6m_{0s}^4}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\lambda_s^k\frac{K_2(kx_s)}{(kx_s)^2}\} \end{aligned}$$

. ...(4)

4.3 Baryon Number Conservation Equation

The net baryon number of the system equals

$$n_B V = \frac{1}{3}(n_q - n_{\overline{q}})V$$

If we restrict our discussions to a unit volume then the baryon density equals

$$n_B = \frac{1}{3}(n_q - n_{\overline{q}}) = \frac{4T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} - \lambda_{\overline{q}}^k e^{-\frac{k\mu_q}{T}}}{k^3}\}$$

for given values of the parameters. Thus, if the initial baryon number density, temperature and non-equilibrium fugacities be known (as from models like HIJING or SSPC or likewise) then we can iterate to get initial value of the quark chemical potential.

The baryon number conservation equation gives

$$\partial_\mu \ (n_B u^\mu) = \frac{\partial n_B}{\partial \tau} + \frac{n_B}{\tau} = 0 \qquad ...(5)$$

from which we can obtain expression for $\dot{\mu}_q$, which gives the rate of change of the chemical potential with time. If we substitute for the number densities and solve for $\dot{\mu}_q$ we get

$$\dot{\mu}_q = \dot{\lambda}_q B_1 + \dot{\lambda}_{\overline{q}} B_2 + \dot{T} B_3 + B_4 \qquad ...(6)$$

where

$$B_1 = -\frac{T}{\triangle}\sum_{k=1}^{\infty}\frac{(-1)^{k-1}}{k^2}\lambda_q^{k-1} e^{\frac{k\mu_q}{T}}$$

$$B_2 = \frac{T}{\triangle}\sum_{k=1}^{\infty}\frac{(-1)^{k-1}}{k^2}\lambda_{\overline{q}}^{k-1} e^{-\frac{k\mu_q}{T}}$$

$$B_3 = -\frac{3}{\triangle}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} + \lambda_{\overline{q}}^k e^{-\frac{k\mu_q}{T}}}{k^3}\} - \frac{\mu_q}{T}$$

and

$$B_4 = -\frac{T}{\tau\triangle}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} - \lambda_{\overline{q}}^k e^{-\frac{k\mu_q}{T}}}{k^3}\}$$

with

$$\triangle = \Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} + \lambda_{\overline{q}}^k e^{-\frac{k\mu_q}{T}}}{k^2}\}$$

We shall see the effect of having these nonzero values very shortly.

4.4 Quark & Anti-Quark Number Density Evolution Equation

Using standard techniques we can obtain the fermionic number density evolution equations. The marked difference from the zero chemical potential case is that due to the nonzero values of the coefficients of the Baryon Number Conservation equation, the massless quark and antiquark number density evolution equations both depend on either of the two growth rates. Let us put forward the two Number Density Evolution Equations. We include a new process called the quark flavour changing process (qfcp) into our framework.

4.4.1 Massless Quark Number Density Evolution Equation

If the Quark Number Density Evolution Equation be explicitly worked out it takes the form

$$\dot{\lambda}_q Q_1 + \dot{\lambda}_{\overline{q}} Q_2 + \dot{T} Q_3 + Q_4 = 0 \qquad \ldots(7)$$

where

$$Q_1 = \frac{12T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^{k-1}e^{\frac{k\mu_q}{T}} + \lambda_q^k e^{\frac{k\mu_q}{T}}\frac{B_1}{T}}{k^2}\}$$

$$Q_2 = \frac{12T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_q^k e^{\frac{k\mu_q}{T}}\frac{B_2}{T}}{k^2}$$

$$Q_3 = \frac{36T^2}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_q^{k-1}e^{\frac{k\mu_q}{T}}}{k^3} + \frac{12T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_q^k e^{\frac{k\mu_q}{T}}\frac{B_3}{T}}{k^2}$$

and

$$Q_4 = \frac{12T^3}{\tau\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_q^k e^{\frac{k\mu_q}{T}}}{k^3} + \frac{12T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_q^k e^{\frac{k\mu_q}{T}}\frac{B_4}{T}}{k^2} \quad - SQ$$

where

$$SQ = (R_{gg\rightarrow q\overline{q}} - R_{q\overline{q}\rightarrow gg}) + (R_{s\overline{s}\rightarrow q\overline{q}} - R_{q\overline{q}\rightarrow s\overline{s}})$$

4.4.2 Massless Anti-Quark Number Density Evolution Equation

For anti-quarks the equations are more or less identical as above with $\lambda_q \rightarrow \lambda_{\bar{q}}$ and $\mu_q \rightarrow -\mu_q$, except for the modifications in the contributions from B_i . If the Anti-Quark Number Density Evolution Equation be explicitly worked out it takes the form

$$\dot{\lambda}_q AQ_1 + \dot{\lambda}_{\overline{q}} AQ_2 + \dot{T} AQ_3 + AQ_4 = 0 \qquad \ldots(8)$$

where

$$AQ_1 = -\frac{12T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_{\bar{q}}^{k}e^{-\frac{k\mu_q}{T}}\frac{B_1}{T}}{k^2}$$

$$AQ_2 = \tfrac{12T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\tfrac{\lambda_{\bar{q}}^{k-1}e^{-\frac{k\mu_q}{T}}-\lambda_{\bar{q}}^{k}e^{-\frac{k\mu_q}{T}}\frac{B_2}{T}}{k^2}\}$$

$$AQ_3 = \frac{36T^2}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_{\bar{q}}^{k-1}e^{-\frac{k\mu_q}{T}}}{k^3} - \frac{12T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_{\bar{q}}^{k}e^{-\frac{k\mu_q}{T}}\frac{B_3}{T}}{k^2}$$

$$AQ_4 = \frac{12T^3}{\tau\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_{\bar{q}}^{k}e^{-\frac{k\mu_q}{T}}}{k^3} - \frac{12T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{\lambda_{\bar{q}}^{k}e^{-\frac{k\mu_q}{T}}\frac{B_4}{T}}{k^2} - SaQ$$

where

$$SaQ = (R_{gg\to q\overline{q}} - R_{q\overline{q}\to gg}) + (R_{s\overline{s}\to q\overline{q}} - R_{q\overline{q}\to s\overline{s}}) = SQ$$

4.4.3 Massive Strange Quark Number Density Evolution Equation

For massive strange quark the evolution equation is given by

$$\dot{\lambda}_s S_1 + \dot{T} S_2 + S_3 = 0 \qquad ...(9)$$

where

$$S_1 = \frac{\sum_{k=1}^{\infty}(-1)^{k-1}k\lambda_s^{k-1}\frac{\dot{K}_2(kx_s)}{(kx_s)}}{\sum_{k=1}^{\infty}(-1)^{k-1}\lambda_s^{k}\frac{\dot{K}_2(kx_s)}{(kx_s)}}$$

$$S_2 = \frac{3}{T}\frac{\sum_{k=1}^{\infty}(-1)^{k-1}\lambda_s^{k}\{\frac{\dot{K}_2(kx_s)}{(kx_s)} + \frac{1}{3}K_1(kx_s)\}}{\sum_{k=1}^{\infty}(-1)^{k-1}\lambda_s^{k}\frac{\dot{K}_2(kx_s)}{(kx_s)}}$$

$S_3 = \frac{1}{\tau} - SQs$ .... (11c) with

$$SQs = \{(R_{gg\to s\overline{s}} - R_{s\overline{s}\to gg})/n_s\} + 2\{(R_{q\overline{q}\to s\overline{s}} - R_{s\overline{s}\to q\overline{q}})/n_s\}$$

4.5 Energy-Momentum Conservation Equation

The conservation of the energy-momentum tensor , which we term as Bjorken's equation for future references,gives

$$\frac{d\varepsilon}{d\tau} + \frac{\varepsilon + p}{\tau} = 0 \qquad ...(10)$$

Inserting the full expressions of energy density and pressure we obtain

$$\dot{T}f_6 + \dot{\lambda}_g f_7 + \dot{\lambda}_q f_8 + \dot{\lambda}_{\overline{q}} f_9 + \dot{\lambda}_s f_s + f_{10} = 0 \qquad ...(11)$$

where

$$f_6 = \frac{192T^3}{\pi^2}\sum_{k=1}^{\infty}\frac{\lambda_g^k}{k^4} + \frac{36T^4}{\pi^2}(\frac{B_3}{T} - \frac{\mu_q}{T^2})\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} - \lambda_{\bar{q}}^k e^{-\frac{k\mu_q}{T}}}{k^3}\}+$$

$$\frac{144T^3}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} + \lambda_{\bar{q}}^k e^{-\frac{k\mu_q}{T}}}{k^4}\} + \frac{72m_{0s}^4}{\pi^2 T}\Sigma_{k=1}^{\infty}(-1)^{k-1}\lambda_s^k\{\frac{K_2(kx_s)}{(kx_s)}+$$

$$+\frac{5K_1(kx_s)}{12(kx_s)} + \frac{K_0(kx_s)}{12}\}$$

$$f_7 = \frac{48T^4}{\pi^2}\sum_{k=1}^{\infty}\frac{\lambda_g^{k-1}}{k^3}$$

$$f_8 = \frac{18T^4}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^{k-1} e^{\frac{k\mu_q}{T}}}{k^2}\} + \frac{18T^4}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{(\lambda_q^{k-1} e^{\frac{k\mu_q}{T}} - \lambda_q^k e^{\frac{k\mu_q}{T}})\frac{B_1}{T}}{k^3}$$

$$f_9 = \frac{18T^4}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_{\bar{q}}^{k-1} e^{-\frac{k\mu_q}{T}}}{k^2}\} + \frac{18T^4}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\frac{(\lambda_q^{k-1} e^{\frac{k\mu_q}{T}} - \lambda_q^k e^{\frac{k\mu_q}{T}})\frac{B_2}{T}}{k^3}$$

$$f_s = \frac{6m_{0s}^4}{\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}k.\lambda_s^{k-1}\{\frac{3K_2(kx_s)}{(kx_s)} + \frac{K_1(kx_s)}{(kx_s)}\}$$

$$f_{10} = \frac{64T^4}{\tau\pi^2}\sum_{k=1}^{\infty}\frac{\lambda_g^k}{k^4} + \frac{48T^4}{\tau\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} + \lambda_{\bar{q}}^k e^{-\frac{k\mu_q}{T}}}{k^4}\}+$$

$$\frac{6m_{0s}^4}{\tau\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\lambda_s^k\{\frac{4K_2(kx_s)}{(kx_s)}+\frac{K_1(kx_s)}{(kx_s)}\}+$$

$$\frac{36T^3}{\tau\pi^2}\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} - \lambda_{\bar{q}}^k e^{-\frac{k\mu_q}{T}}}{k^3}\}B_4$$

5 Changing Distribution Functions

In order to reduce computational complications to a low degree we make certain approximations.We redefine the non-equilibrium distribution functions and shift from the Juttner distribution function to Modified Fermi-Dirac type for the quarks and antiquarks in confirmation with earlier works [4,5]. Needless to say, this would surely limit the applicability of the present study. However, it is believed that the basic nature of partonic fugacity variations would remain unaltered.Since we adopt certain approximations for reasons as mentioned earlier, we refrain from giving the explicit forms of the reaction rates presently and hold back the expressions thereof for now.

As stated, we adopt the following approximations:

We replace the original distribution functions by the following:

$$
\begin{aligned}
&f_g = \frac{\lambda_g}{e^{\varepsilon/T} - 1}\\
&f_{q(\overline{q})} = \frac{\lambda_{q(\overline{q})} e^{\pm\frac{\mu_q}{T}}}{e^{\varepsilon/T} + 1} \qquad\qquad\qquad\qquad \text{...(12)}\\
&f_s = f_{\overline{s}} = \frac{\lambda_s}{e^{\varepsilon/T} + 1}
\end{aligned}
$$

Of these, the last two are the Modified Fermi-Dirac (MFD) type Distribution functions as was used in [4,5] .

As for the quarks, we take
i) For s quark we take mass~ 150 MeV.
ii) For lighter quarks since mass is much too less than the strange quark mass, we treat the mass of lighter flavours as zero.
iii) Since the lighter quarks are treated to have the same mass, we neglect $u\overline{u} \rightleftharpoons d\overline{d}$ type of reactions as they would be like elastic scattering processes leading to no net particle production.

5.1 Basic Thermodynamic Quantities

Using the techniques of statistical mechanics, we can obtain the number density, energy density and pressure as follows:

$$
\begin{aligned}
&n = \frac{16\zeta(3)T^3}{\pi^2}\lambda_g + \frac{9\zeta(3)n_f T^3}{2\pi^2}\{\lambda_q e^{\mu_q/T} + \lambda_{\overline{q}} e^{-\mu_q/T}\} +\\
&\frac{6m_{0s}^3}{\pi^2}\lambda_s \sum_{k=1}^{\infty}(-1)^{k-1}\frac{K_2(kx_s)}{(kx_s)}\\
&\varepsilon = \frac{8\pi^2 T^4}{15}\lambda_g + \frac{7\pi^2 n_f T^4}{40}\{\lambda_q e^{\mu_q/T} + \lambda_{\overline{q}} e^{-\mu_q/T}\} +\\
&\frac{6m_{0s}^4}{\pi^2}\lambda_s \sum_{k=1}^{\infty}(-1)^{k-1}\{\frac{3K_2(kx_s)}{(kx_s)^2} + \frac{K_1(kx_s)}{(kx_s)}\}\\
&p = \frac{8\pi^2 T^4}{45}\lambda_g + \frac{7\pi^2 n_f T^4}{120}\{\lambda_q e^{\mu_q/T} + \lambda_{\overline{q}} e^{-\mu_q/T}\} +\\
&\frac{6m_{0s}^4}{\pi^2}\lambda_s \sum_{k=1}^{\infty}(-1)^{k-1}\frac{K_2(kx_s)}{(kx_s)^2}\\
&with \;\; x_s = m_s/T
\end{aligned}
$$

...(13)

Using standard techniques we can obtain the quark and anti-quark number density evolution equations. As for the Juttner distribution case, the equations are coupled via the Baryon Number Conservation Equation coefficients B_i .

5.2 Baryon Number Conservation Equation

Using the modified distribution functions we can obtain the coefficients B_i . We obtain

$\dot{\mu}_q = \dot{\lambda}_q B_1 + \dot{\lambda}_{\overline{q}} B_2 + \dot{T} B_3 + B_4$ ...(14)

where

$$B_1 = -\frac{T e^{\frac{\mu_q}{T}}}{\Delta_1}$$

$$B_2 = \frac{T e^{-\frac{\mu_q}{T}}}{\Delta_1}$$

$$B_3 = \frac{\mu_q}{T} - \frac{\Delta_2}{\Delta_1}$$

$$B_4 = -\frac{T}{\tau}\frac{\Delta_2}{\Delta_1}$$

where

$$\Delta_1 = \lambda_i e^{\frac{\mu_i}{T}} + \lambda_{\overline{i}} e^{-\frac{\mu_i}{T}}$$

$$\Delta_2 = \lambda_i e^{\frac{\mu_i}{T}} - \lambda_{\overline{i}} e^{-\frac{\mu_i}{T}}$$

5.3 Quark and Anti-Quark Number Density Evolution Equations

Just as in the case of the full Juttner distribution, here too the quark and anti-quark number density evolution equations are coupled via the coefficients B_i .

5.3.1 Massless Quark Number Density Evolution Equation

The equation is given by

$\dot{\lambda}_q Q_1 + \dot{\lambda}_{\overline{q}} Q_2 + \dot{T} Q_3 + Q_4 = 0$ ...(15)

where

$$Q_1 = \frac{1}{\lambda_q} + \frac{B_1}{T}$$

$$Q_2 = \frac{B_2}{T}$$

$$Q_3 = \frac{3}{T} + \frac{B_3}{T} - \frac{\mu_q}{T^2}$$

$$Q_4 = \frac{1}{\tau} + \frac{B_4}{T} - SQ \text{ with}$$
$$SQ = \{(R_{gg\to q\overline{q}} - R_{q\overline{q}\to gg})/n_q\} - \{(R_{q\overline{q}\to s\overline{s}} - R_{s\overline{s}\to q\overline{q}})/n_q\}$$

5.3.2 Massless Anti-Quark Number Density Evolution Equation

The equation is given by

$$\dot{\lambda}_q Q_1 + \dot{\lambda}_{\overline{q}} Q_2 + \dot{T} Q_3 + Q_4 = 0 \qquad ...(16)$$

$$AQ_1 = -\frac{B_1}{T}$$
$$AQ_2 = \frac{1}{\lambda_{\overline{q}}} - \frac{B_2}{T}$$
$$AQ_3 = \frac{3}{T} - \frac{B_3}{T} + \frac{\mu_q}{T^2}$$
$$AQ_4 = \frac{1}{\tau} - \frac{B_4}{T} - SaQ \quad \text{with}$$
$$SaQ = \{(R_{gg\to q\overline{q}} - R_{q\overline{q}\to gg})/n_{\overline{q}}\} - \{(R_{q\overline{q}\to s\overline{s}} - R_{s\overline{s}\to q\overline{q}})/n_{\overline{q}}\}$$

5.3.3 Massive Strange Quark Number Density Evolution Equation

For massive strange quark the evolution equation is given by

$$\dot{\lambda}_s S_1 + \dot{T} S_2 + S_3 = 0 \qquad ...(17)$$

where

$$S_1 = \frac{1}{\dot{\lambda}_s}$$

$$S_2 = \frac{3}{T} \frac{\sum_{k=1}^{\infty} (-1)^{k-1} \{ \frac{K_2(kx_s)}{(kx_s)} + \frac{1}{3} K_1(kx_s) \}}{\sum_{k=1}^{\infty} (-1)^{k-1} \lambda_s^k \frac{K_2(kx_s)}{(kx_s)}}$$

$$S_3 = \frac{1}{\tau} - SQs \text{ with}$$
$$SQs = \{(R_{gg\to s\overline{s}} - R_{s\overline{s}\to gg})/n_s\} + 2\{(R_{q\overline{q}\to s\overline{s}} - R_{s\overline{s}\to q\overline{q}})/n_s\}$$

5.4 Bjorken's Equation

From Bjorken's equation we get on substituting for energy and momentum

$$\dot{T} f_6 + \dot{\lambda}_g f_7 + \dot{\lambda}_q f_8 + \dot{\lambda}_{\overline{q}} f_9 + \dot{\lambda}_s f_s + f_{10} = 0 \qquad ...(18)$$

where

$$f_6 = \frac{32\pi^2}{15}\lambda_g T^3 +$$

$$\frac{7\pi^2 n_f}{40} T^4 (\lambda_q e^{\frac{\mu_q}{T}} - \lambda_{\overline{q}} e^{-\frac{\mu_q}{T}})(\frac{B_3}{T} - \frac{\mu_q}{T^2}) + \frac{28\pi^2 n_f}{40} T^3 (\lambda_q e^{\frac{\mu_q}{T}} + \lambda_{\overline{q}} e^{-\frac{\mu_q}{T}}) +$$

$$\frac{72 m_{0s}^4}{\pi^2 T}\lambda_s \sum_{k=1}^{infty} (-1)^{k-1} \{ \frac{K_2(kx_s)}{(kx_s)^2} + \frac{5K_1(kx_s)}{12(kx_s)} + \frac{K_0(kx_s)}{12} \}$$

$$f_7 = \frac{8\pi^2}{15} T^4$$

$$f_s = \frac{6 m_{0s}^4}{\pi^2} \sum_{k=1}^{\infty} (-1)^{k-1} \{ \frac{3K_2(kx_s)}{(kx_s)^2} + \frac{K_1(kx_s)}{(kx_s)} \}$$

$$f_8 = \frac{7\pi^2 n_f}{40} \{ e^{\frac{\mu_q}{T}} + (\lambda_q e^{\frac{\mu_q}{T}} - \lambda_{\overline{q}} e^{-\frac{\mu_q}{T}}) \frac{B_1}{T} \}$$

$$f_9 = \frac{7\pi^2 n_f}{40} \{ e^{-\frac{\mu_q}{T}} + (\lambda_q e^{\frac{\mu_q}{T}} - \lambda_{\overline{q}} e^{-\frac{\mu_q}{T}}) \frac{B_2}{T} \}$$

$$f_{10} = \frac{32\pi^2}{45\tau}\lambda_g T^4 + \frac{6 m_{0s}^4}{\pi^2 \tau}\lambda_s \Sigma_{k=1}^{\infty} (-1)^{k-1} \{ \frac{4K_2(kx_i)}{(kx_i)^2} + \frac{K_1(kx_i)}{(kx_i)} \}$$

$$+ \frac{28\pi^2 n_f}{120} T^4 (\lambda_q e^{\frac{\mu_q}{T}} + \lambda_{\overline{q}} e^{-\frac{\mu_{qi}}{T}}) + \frac{7\pi^2 n_f}{40} T^4 (\lambda_q e^{\frac{\mu_q}{T}} - \lambda_{\overline{q}} e^{-\frac{\mu_q}{T}}) \frac{B_4}{T}$$

Now,it is to be noted as the quark and anti-quark number density evolution equations and the baryon number conservation equation are mathematically very simply connected, so these two evolution equations linked by the four B coefficients along with the energy-momentum conservation equation with inputs from the gluon and strange quark number density evolution equations do not constitute a set of solvable equations as the system determinant vanishes. To make the system solvable, we expand the exponential term in the numerators of the first two B coefficients and thereby introduce some 'assymetry'. Strictly speaking, this however would mean not exact baryon conservation.

5.5 The Partonic Reaction Rates

5.5.1 The Gluon Multiplication Rate R_3

The gluon multiplication rate has been calculated by Xiong et. al. [10]. The rate of the reaction $gg \to (n-2)g$ is given. By explicitly calculating the ma-

trix element [11] (summed over all the final states and averaged over all initial states) we can obtain the gluon multiplication rate.
However, to avoid the huge calculations of evaluating 25 Feynman diagrams [11] involved, we fall back on previous results that earlier workers had used . We postulate that the gluon multiplication rate depends on the chemical potential via the Debye Screening mass.
The Debye Screening mass suitable for a multicomponent chemically non-equilibrated parton plasma is given by [12]

$$m_D^2 = \frac{2g^2}{\pi^2}\int dkk[N_c f_g + \textstyle\sum_i f_i] \qquad ...(19)$$

where the sum runs over all flavours i, while N_c gives the number of colours.
To accomodate for antiquarks and remembering that our number of flavours is 3 and not 6, we propose the following modification:

$$m_D^2 = \frac{2g^2}{\pi^2}\int dkk[3f_g + \frac{1}{2}\ \textstyle\sum_{i=u,d,s}(f_i + f_{\bar{i}})] \qquad ...(20)$$

Using standard techniques , we can obtain the following results for the mean free path:

$$\lambda_f^{-1} = n_g \textstyle\int dq_\perp^2 \frac{d\sigma_{el}^{gg}}{dq_\perp^2} = \frac{9n_g\pi\alpha_s^2}{2m_D^2(1+\frac{2}{9}\frac{m_D^2}{T^2})} \qquad ...(21)$$

which for zero chemical potential reduces to the well known result:

$$\lambda_f^{-1} = \frac{9}{8}a_1\alpha_s T\frac{1}{1+8\pi\alpha_s\lambda_g/9}$$

Using methods of our previous chapter we get on integrating the modified differential cross section and recalling the definition of $R_3 = \frac{1}{2}\sigma_3 n_g$ the required rate as

$$\frac{R_3}{T} = \frac{27\alpha_s^3}{2}\lambda_f^2 n_g I(\lambda_g) \qquad ...(22)$$

where

$$I(\lambda_g) = \int_1^{\sqrt{s}\lambda_f} dx \int_0^{\frac{s}{4m_D^2}} dz \frac{z}{(1+z)^2}[\frac{cosh^{-1}\sqrt{x}}{x\sqrt{[x+(1+z)x_D]^2 - 4xzx_D}} + \frac{1}{s\lambda_f^2}\frac{cosh^{-1}\sqrt{x}}{\sqrt{[1+x(1+z)y_D]^2 - 4xzy_D}}]$$

with

$$x_D = m_D^2\lambda_f$$

$$y_D = \frac{m_D^2}{s}$$

5.5.2 The Quark Anti-Quark pair production rate

We have parton production rates in the RHS of the number density evolution equations.Let us evaluate the quark-antiquark pair production reaction rate R_{2g} . We have [8]

$$R_{2g} = R_{gain}^{gg} - R_{loss}^{gg} \qquad ...(23)$$

where

$$R^{gg}_{gain} = \int \frac{d^3p_1}{(2\pi)^3 2E_1} \int \frac{d^3p_2}{(2\pi)^3 2E_2} \int \frac{d^3p_3}{(2\pi)^3 2E_3} \int \frac{d^3p_4}{(2\pi)^3 2E_4} (2\pi)^4$$

$$\delta^4(p_1 + p_2 - p_3 - p_4) \Sigma |M_{gg \to i\bar{i}}|^2 f_g(p_1) f_g(p_2)(1 - f_q(p_3))(1 - f_{\bar{q}}(p_4)) \qquad ...(23a)$$

and

$$R^{gg}_{loss} = \int \frac{d^3p_1}{(2\pi)^3 2E_1} \int \frac{d^3p_2}{(2\pi)^3 2E_2} \int \frac{d^3p_3}{(2\pi)^3 2E_3} \int \frac{d^3p_4}{(2\pi)^3 2E_4} (2\pi)^4$$

$$\delta^4(p_1 + p_2 - p_3 - p_4) \Sigma |M_{gg \to i\bar{i}}|^2 (1 + f_g(p_1))(1 + f_g(p_2)) f_q(p_3) f_{\bar{q}}(p_4) \qquad ...(23b)$$

Following [8], we can say that there are three topologically distinct Feynmann diagrams that contribute towards the quark-antiquark pair production process. Evaluating them performing traces and finally adding them up we can find the net squared matrix element. We basically follow the lines of [8]. Transforming variables as

$$q = p_1 + p_2$$

$$p = \frac{1}{2}(p_1 - p_2)$$

$$q' = p_3 + p_4$$

$$p' = \frac{1}{2}(p_3 - p_4)$$

with restrictions

$$q_0 > 2m_i$$

$$s = q_0^2 - |\overrightarrow{\mathbf{q}}|^2 \geq 4m_i^2$$

$$p_0^2 \leq \frac{q^2}{4}$$

$$p_0' . p_0' \leq \frac{q^2}{4}(1 - \frac{4m_i^2}{s})$$

and transforming the three dimensional integrals to four dimensional integrals using

$$\int \frac{d^3p_i}{2E_i} = \int d^4p_i \delta(p^2 - m_i^2)$$

with the new set of variables

$$q_0 = -T ln v + 2m_i$$

$$q^{\frac{1}{2}} = (q_0^2 - 4m_i^2)^{\frac{1}{2}} u$$

$$p_0 = \frac{q}{2}(1 - \frac{4m_i^2}{s})^{\frac{1}{2}} x$$

$$p_0' = \frac{q}{2} y$$

we arrive at the rate

$$R_{2g} = \frac{\alpha_s^2}{2\pi^3} T \int_0^1 du \int_0^1 dv \int_0^1 dx \int_0^1 dy \frac{u^2}{v} (1 - \frac{4m_i^2}{s})^{\frac{1}{2}} (q_0^2 - 4m_i^2)^{\frac{3}{2}} f_{Quarks} f_{phase1} \quad ...(24)$$

where

$$f_{Quarks} = f_g(\frac{q_0}{2} + p_0) f_g(\frac{q_0}{2} - p_0)(1 - f_q(\frac{q_0}{2} + p_0'))(1 - f_{\bar{q}}(\frac{q_0}{2} - p_0')) -$$
$$(1 + f_g(\tfrac{q_0}{2} + p_0))(1 + f_g(\tfrac{q_0}{2} - p_0)) f_q(\tfrac{q_0}{2} + p_0') f_{\bar{q}}(\tfrac{q_0}{2} - p_0')$$

and

$$f_{phase1} = A + B[\frac{1}{K_+} + \frac{1}{K_-}] + C[\frac{\triangle_+}{K_+^3} + \frac{\triangle_-}{K_+^3}]$$

with

$$A = 3[1-[1-\frac{4m_i^2}{s}][\frac{(1-x^2)(1-y^2)}{2}+x^2y^2]]-\frac{34}{3}-24\frac{m_i^2}{s}$$
$$B = \frac{16}{3}[1+\frac{4m_i^2}{s}+\frac{m_i^4}{s^2}]$$
$$C = -\frac{128}{3}\frac{m_i^4}{s^2}$$
$$K_{\pm} = [1-[1-\frac{4m_i^2}{s}][(1-x^2-y^2)\pm 2[1-\frac{4m_i^2}{s}]^{\frac{1}{2}}xy]^{\frac{1}{2}}$$

5.5.3 The Quark flavour changing rate

For the quark flavour changing process we have [8]

$$R_{qg} = R^{q\overline{q}} - R^{q\overline{q}}_{loss} \qquad ...(25)$$

where

$$R^{q\overline{q}}_{gain} = \int \frac{d^3p_1}{(2\pi)^3 2E_1} \int \frac{d^3p_2}{(2\pi)^3 2E_2} \int \frac{d^3p_3}{(2\pi)^3 2E_3} \int \frac{d^3p_4}{(2\pi)^3 2E_4} (2\pi)^4$$
$$\delta(p_1+p_2-p_3-p_4)\Sigma|M_{s\overline{s}\rightarrow q\overline{q}}|^2 f_q(p_1)f_{\overline{q}}(p_2)(1-f_s(p_3))(1-f_{\overline{s}}(p_4)) \qquad ...(25a)$$

and

$$R^{q\overline{q}}_{loss} = \int \frac{d^3p_1}{(2\pi)^3 2E_1} \int \frac{d^3p_2}{(2\pi)^3 2E_2} \int \frac{d^3p_3}{(2\pi)^3 2E_3} \int \frac{d^3p_4}{(2\pi)^3 2E_4} (2\pi)^4$$
$$\delta^4(p_1+p_2-p_3-p_4)\Sigma|M_{s\overline{s}\rightarrow q\overline{q}}|^2 (1-f_q(p_1))(1-f_{\overline{q}}(p_2))f_s(p_3)f_{\bar{s}}(p_4) \qquad ...(25b)$$

Following [8], we can say that there is only one type of topologically distinct Feynmann diagram that contributes towards the quark flavour changing process or the strange quark pair production process. Evaluating it, performing trace calculations we can find the squared matrix element. We basically follow the lines of earlier works. Transforming variables as in the case before and performing identical operations we can get the rate as

$$R_{qg} = \frac{\alpha_s^2}{2\pi^3}T\int_0^1 du \int_0^1 dv \int_0^1 dx \int_0^1 dy \frac{u^2}{v}(1-\frac{4m_i^2}{s})^{\frac{1}{2}}(q_0^2-4m_i^2)^{\frac{3}{2}} f_{Strange}f_{phase2} \qquad ...(26)$$

with

$$f_{Quarks} = f_q(\frac{q_0}{2}+p_0)f_{\overline{q}}(\frac{q_0}{2}-p_0)(1-f_s(\frac{q_0}{2}+p_0'))(1-f_s(\frac{q_0}{2}-p_0'))-$$
$$(1+f_s(\frac{q_0}{2}+p_0))(1+f_s(\frac{q_0}{2}-p_0))f_q(\frac{q_0}{2}+p_0')f_{\overline{q}}(\frac{q_0}{2}-p_0')$$
$$f_{phase2} = [1+[1-\frac{4m_i^2}{s}][\frac{(1-x^2)(1-y^2)}{2}+x^2y^2]]+\frac{4m_i^2}{s}$$

6 Results

The initial Conditions from the Self-Screened-Parton-Cascade (SSPC) Model are as follows:

	$\tau_i (fm)$	$T(GeV)$	λ_g	λ_q		λ_s
RHIC	0.25	0.668	0.34	0.064		0.032
LHC	0.25	1.02	0.43	0.086		0.043

In absence of known value of initial baryon number density at the point when thermal equilibrium is attained (after which point the system evolves according to the laws of hydrodynamics) we perform a study with varying initial conditions. We vary the initial baryon number density between 0.11 and 0.21/fm^3 while we change the initial quark to antiquark fugacity ratio between 1.1 and 1.9. Needless to say, most set of values would be of academic interest only and not physically realisable. But, this could lead to a better understanding, it is believed.

We study comparatively the outputs with given initial baryon number density and initial light quark to anti-quark fugacity ratio. We arrive at the following conclusions :

1. The nature of variations of the physical quantities are more or less in the line of earlier works. The variations of Temperature, Chemical Potential and Non equilibrium Fugacities for RHIC and LHC initial conditions are shown in figures 1 and 2 respectively. The decaying curves give temperature variations while the positive rising curves are sequentially (from top) for gluon, light quark, light anti-quark and strange quark fugacity variations. Plots for both quark flavour changing process included and excluded cases are shown. For identification please see point 4 below.As representative values we take baryon number density 0.15 and initial fugacity ratio 1.5.

We observe that
i) As expected, the temperature falls with time while the non-equilibrium fugacities increase.
ii) Contrary to the Juttner case, the chemical potential remains negative all along and as expected approaches zero as the system equilibrates.
iii) The QGP,as expected remains to be gluon dominated.
iv) For a system at higher chemical potential, the temperature falls at a slower rate signifying lesser amount of energy expenditure to create partons which shows up in the slower rise of all partons except for the light anti-quark, which shows a larger growth rate due to the presence of the exponentiated chemical potential.

2. For a given initial ratio, except for the light quarks and anti-quarks, the output does not depend much on the initial baryon number density. For the

light quarks and antiquarks this variation is due to the presence of the exponentiated non-zero chemical potentials.

3. For a given initial baryon number density we can recast the equation for baryon number density in the form

$$e^{\mu_q/T} = \frac{C}{2\lambda_q} + \sqrt{\frac{1}{D} + \frac{C^2}{4\lambda_q^2}}$$

, ...(27)

where C is a constant for a given temperature, light quark fugacity and given baryon number density. Here D is the light quark to antiquark initial fugacity ratio. For a fixed light quark fugacity and temperature, as D falls clearly the RHS of the above equation increases which indicates a rise in the chemical potential. Again for a system of higher chemical potential, the temperature has to drop at a slower rate due to the constraint imposed. Hence we observe that

i) The temperature falls at a slower rate for a smaller value of the light quark to antiquark initial fugacity ratio.
ii) As the temperature falls at a slower rate, it would imply a lesser expenditure of energy to produce particles in general . This would show up in the slower rise of all fugacity values except the light antiquark, which would show a higher growth rate. This is due to the exponentiated chemical potential part.

4. For inclusion of the quark-flavour-changing process we observe the following:

Due to the additional production of s-quarks from massless quarks via qfcp, we see an additional increase in fugacity of the strange quark while, the rate of equilibration falls for non-strange fermions .

5. For some initial baryon number density and known values of non-equilibrium fugacities and temperature switching distribution functions would mean a shift in the value of the light quark chemical potential, which is found by iteration. Since everything else remains the same, to fit the baryon number density, clearly the chemical potential is expected to 'pay the price'. This is exactly what happens. For the MFD case, the chemical potential assumes a value much smaller (-ve and also magnitudinally an order smaller) than that for the Juttner case (in which case it is +ve). In the result for the number density using the Juttner distributions we had an expression like $\Sigma_{k=1}^{\infty}(-1)^{k-1}\{\frac{\lambda_q^k e^{\frac{k\mu_q}{T}} + \lambda_{\bar{q}}^k e^{-\frac{k\mu_q}{T}}}{k^3}\}$. If we expand the exponential with an objective of suitable truncation, keeping a 'k' factor in the numerator of the exponent with a much larger value of the chemical potential, it would make the mathematics still more complex as it would be required to consider many more number of terms for the same accuracy. Therefore, we had to adopt the MFD type of distribution functions (

with k=1 in the exponent), where, as required, we can easily use a truncated expansion of the exponential term.

7 Acknowledgement

The author gratefully acknowledges helpful discussions with Prof. Bikash Sinha and Prof.Binayak Dutta Roy as also valuable e-mail clarifications and encouragement from Profs. Z.J.He and Y.G.Ma during the initial stages of the present work.

8 References

1. N.Hammon et. al, Phys. Rev. C61, 014901 (1999)
2. STAR Collaboration, Phys. Lett. B567 (2003)167
3. BRAHMS Collaboration, Phys. Lett. B607 (2005)42
4. D.Dutta, A.K.Mohanty, K.Kumar, R.K.Choudhury,Phys.Rev.C60(1999)014905
5. D.Dutta, A.K.Mohanty, K.Kumar, R.K.Choudhury,Phys.Rev.C61(2000)064911
6. A.Sen, Rev. Bull. Cal. Math. Soc. 11 (1 & 2) 39-44 (2003)
7. Z.J.He, J.L.Long, Y.G.Ma, G.L.Ma B.Liu, Phys. Rev. C69, 034901 (2004)
8. T.Matsui, B.Svetitsky, L.D.McLerran, Phys. Rev. D34, 783 (1986)
9. Introduction to Relativistic Heavy Ion Collisions
Laszo P. Csernai, John Wiley and Sons, 1994
10.Li Xiong et.al., Phys.Rev.C49,2203 (1994)
11.F.A.Berends et.al, Phys. Lett. 103B,124(1981)
12. Fred Cooper et. al., www.arXiv.org/hep-ph/0207370


Figure2: LHC Case
Temperature(GeV),Chemical Potential(GeV) & Fugacity
1.2
1
0.8
0.6
0.4
0.2
0
-0.2
5
10
15
20
25
30
Proper time in fm

Figure1: RHIC Case